



The ibidi Mounting Medium is a glycerol-based mounting solution optimized for ibidi  $\mu$ -Slides,  $\mu$ -Dishes, and  $\mu$ -Plates with adherent cells or tissue sections that are immunofluorescent stained. It contains an inhibitor that retards photobleaching of fluorochromes. ibidi Mounting Medium is ready-to-use.

## Overview

This document is applicable to the following product numbers:

Cat. No.	Product Name
50001	ibidi Mounting Medium: for fluorescence microscopy, 15 ml
50001-4	ibidi Mounting Medium: for fluorescence microscopy, 4x15 ml

## Material

ibidi Mounting Medium is glycerol-based, non-drying and non-hardening. It contains 1,4-diazabicyclo[2.2.2]octane (DABCO) to prevent photobleaching. It is dilutable with water or phosphate buffered saline (PBS) to a maximum dilution of 50%. Diluted ibidi Mounting Medium can lose volume by evaporation after some time. Undiluted ibidi Mounting Medium is volume-stable with a maximum change of volume of approx. 10% depending on the sample storage conditions.

Specifications	
Appearance	Liquid, optically clear
pH	8.5–9.0
Refractive index $n_D$	1.42–1.44
Storage of mounted samples	In the dark at 2–8°C

## Shipping and Storage

Shipping conditions	Ambient
Storage conditions	RT (15–25°C)
Shelf life	Under proper storage conditions as indicated on vial


## Applications

ibidi Mounting Medium is designed for immunohistochemistry, immunofluorescent stainings and fluorescence in situ hybridization (FISH) of adherent cells or tissue sections. It is compatible with plastic and glassware.

## Usage

### Open wells and dishes

After fixation and staining wash thoroughly with PBS or appropriate buffer and place some drops of ibidi Mounting Medium onto the cells using the dropper bottle. The volume depends on the used well or dish diameter. It is recommended filling the entire growth area of the well or dish.



### Channel slides

For channel slides like  $\mu$ -Slide VI<sup>0.4</sup>, press the dropper bottle tightly onto the Luer adapter and inject the Mounting Medium into the empty channel. Small remains of PBS or appropriate buffer inside the channel do not cause problems. It is recommended filling the entire channel.



### Sample storage

Store mounted samples under cool and dark conditions. Keep all samples upright not to cause leakage. Samples can be stored for months and years depending on the stability of antibodies and fluorochromes.

**Ordering Information**

The ibidi Mounting Medium family is available with or without DAPI.

Cat. No.	Description	Amount
50001	<b>ibidi Mounting Medium:</b> for fluorescence microscopy	1 × 15 ml
50001-4	<b>ibidi Mounting Medium:</b> for fluorescence microscopy	4 × 15 ml
50011	<b>ibidi Mounting Medium with DAPI:</b> for fluorescence microscopy	1 × 15 ml
50011-4	<b>ibidi Mounting Medium with DAPI:</b> for fluorescence microscopy	4 × 15 ml

**For research use only!**

Further information can be found at [ibidi.com](http://ibidi.com). For questions and suggestions please contact us by e-mail [info@ibidi.de](mailto:info@ibidi.de) or by telephone +49 (0)89/520 4617 0.

© ibidi GmbH, Lochhamer Schlag 11, 82166 Gräfelfing, Germany.